

**CHEMICAL COMPOSITION AND
AUTHENTICITY OF SELECTED EDIBLE BIRD
NEST IN MALAYSIA AND INDONESIA**

NUR HULWANI BINTI IBRAHIM

© This item is protected by original copyright

UNIVERSITI MALAYSIA PERLIS

2015



**Chemical composition and authenticity of selected
Edible Bird Nest in Malaysia and Indonesia**

by

Nur Hulwani Binti Ibrahim

(1231110762)

A thesis submitted in fulfillment of the requirements for the degree of
Master of Science Bioprocess Engineering

**School of Bioprocess Engineering
UNIVERSITI MALAYSIA PERLIS**

2015

ACKNOWLEDGEMENT

First of all, Alhamdulillah, all praise to Allah. Finally, I managed to complete my master thesis after all the hardships and tribulations during this adventurous journey. I would like to express the deepest appreciation to my main supervisor, Assoc. Prof. Dr Zainab Hamzah, which truly helped the progression and smoothness of my research works. The co-operation and tutelage are much indeed appreciated. I am also grateful towards my co-supervisor, Mr Lee Bong Beng for his vital encouragement and support.

It is also my pleasure to express my gratitude to the staffs of School of Bioprocess Engineering, University Malaysia Perlis (UniMAP), especially the teaching engineers and technicians for their tremendous help in analyses, equipment troubleshooting, and other lab work. Not to forget, my gratitude to Assoc. Prof. Dr. Othman for the support and advice and also Dr Ammar and other CEASTech members for providing sensor instrument for my last part of my thesis.

I would like to extend my heartfelt thanks to Higher Education Ministry of Malaysia for the financial support given under KPT (UPM-COE Swiftlet) (9015-00001) and MyBrain 15 (MyMaster).

Last but not least, I would like to express my utmost gratitude to my beloved parents, Mr Ibrahim and Mrs Zainon and my siblings for their love, patience and continuous support.

Finally, I would like to thank all my postgraduate friends for all the support and cheers, especially to Siti Aminah Mohd Hassan, who always accompany me, especially

when I have to stay all night alone finishing the experiment works and Ainatul Alia Alias, who always help me with the lab works.

© This item is protected by original copyright

TABLE OF CONTENTS

	PAGE
THESIS DECLARATION	i
ACKNOWLEDGEMENT	ii
TABLE OF CONTENTS	iv
LIST OF TABLES	ix
LIST OF FIGURES	xi
LIST OF SYMBOLS	xiii
LIST OF ABBREVIATIONS	xiv
ABSTRAK	xvi
ABSTRACT	xvii
CHAPTER 1 INTRODUCTION	
1.1 Background of Study	1
1.2 Statement of Problem	2
1.2.1 Authenticity of EBN	2
1.2.2 Nitrite and Nitrate Content in EBN	3
1.3 Objective of the Study	3
1.4 Scope of the Study	4
1.5 Chapter Summary	4
CHAPTER 2 LITERATURE REVIEW	
2.1 Edible Bird Nest (EBN)	6

2.2	Swiftlets	8
2.3	Origin of Edible Bird Nest	9
2.4	Quality of Edible Bird Nest	10
2.5	Types of Bird Nest	11
2.5.1	Cave Nest	11
2.5.2	House Nest	12
2.6	Quality Assessment of Edible Bird Nest	13
2.6.1	Authenticity of Edible Bird Nest	13
2.6.2	Composition of Edible Bird Nest	15
2.7	Sialic Acid in Edible Bird Nest	20
2.8	Nitrite and Nitrate Content in Edible Bird Nest	21
2.9	Benefits of Edible Bird Nest	23
2.9.1	Nutritional Value of Edible Bird Nest	23
2.9.2	Medicinal Properties of Edible Bird Nest	25
2.10	Chapter Summary	27

CHAPTER 3 RESEARCH METHODOLOGY

3.1	Introduction	28
3.2	Experimental Procedures	29
3.3	Material	30
3.4	Sample Preparation	30
3.5	Characterization of Edible Bird Nest (EBN)	30

3.6	Proximate Analysis	31
3.6.1	Moisture Analysis	31
3.6.2	Ash Analysis	32
3.6.3	Protein Analysis	32
3.6.3.1	Reagents	32
3.6.3.2	Preparation of standard solution for calibration	33
3.6.3.3	Protein Determination in EBN	34
3.6.4	Fat Analysis	34
3.6.4.1	Reagents and apparatus	34
3.6.4.2	Fat Determination in EBN	35
3.6.5	Carbohydrate Analysis	35
3.6.5.1	Reagents	36
3.6.5.2	Carbohydrate Determination in EBN	36
3.6.5.3	Preparation of stock solution	36
3.6.5.4	Preparation of standard solution for calibration	37
3.7	Sialic Acid Analysis	37
3.7.1	Reagent	37
3.7.2	Preparation of stock solution of sialic acid (N-acetylneuraminic acid)	38
3.7.3	Preparation of standard solution for calibration	38
3.7.4	Sialic Acid Determination in EBN	38
3.8	Nitrite and Nitrate Analysis	39
3.8.1	Reagents and apparatus	39

3.8.2	Preparation of standard solution for calibration	39
3.8.3	Nitrate and Nitrite Determination in EBN	40
3.8.4	E-Tongue Measurement	40
3.8.5	Recovery Test	41
3.9	Elemental Analysis	42
3.9.1	Reagent	42
3.9.2	Elemental Determination in EBN	43
3.10	Statistical Analysis	43
3.11	Chapter Summary	44
CHAPTER 4 RESULTS AND DISCUSSIONS		
4.1	Characterization of Edible Bird Nest	45
4.1.1	Analysis of Chemical Compounds and Functional Groups in Pure and Adulterated EBN Using Fourier Transform Infrared (FTIR) Spectroscopy	45
4.2	Moisture Content in EBN Samples	52
4.3	Ash Content in EBN Samples	54
4.4	Protein Content in EBN Samples	56
4.5	Fat Content in EBN Samples	59
4.6	Carbohydrate Content in EBN Samples	61
4.7	Sialic Acid Content in EBN Samples	63
4.8	Determination of Nitrite/Nitrate Content in EBN Samples	66
4.9	Elemental Concentration in EBN Samples	74

CHAPTER 5 CONCLUSIONS & RECOMMENDATIONS

5.1	Conclusions	80
5.2	Recommendations	82

REFERENCES

APPENDIX A	Samples and equipments used in the study	89
APPENDIX B	Characterization of Edible Bird Nest (EBN)	91
APPENDIX C	Calibration curves	96
APPENDIX D	Compositional Analysis of EBN	99
APPENDIX E	JMP Pro 10 Data Sheet	109

LIST OF AWARDS AND PUBLICATIONS

119

LIST OF TABLES

NO.		PAGE
2.1	Protein, carbohydrate and sialic acid content in different types of EBN samples	16
2.2	Crude protein, mineral and sialic acid content in unprocessed and processed EBN	17
2.3	Mineral contents in different types of EBN samples	18
2.4	Nitrite and nitrate content in raw unclean and clean edible bird nest (EBN) from genus <i>Aerodramus</i>	22
4.1	Functional groups of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	48
4.2	Matched Pair t-test to compare the percentage of moisture in raw unclean and clean EBN	53
4.3	Matched Pair t-test to compare the percentage of ash in raw unclean and clean EBN	55
4.4	Matched Pair t-test to compare the percentage of protein in raw unclean and clean EBN	58
4.5	Matched Pair t-test to compare the percentage of fat in raw unclean and clean EBN	60
4.6	Matched Pair t-test to compare the percentage of carbohydrate in raw unclean and clean EBN	62
4.7	Matched Pair t-test to compare the percentage of sialic acid in raw unclean and clean EBN	65
4.8	Nitrite content on Perlis EBN using ion chromatography and sensor techniques	69
4.9	Recovery study performed by adding standard solution of nitrite to Perlis EBN samples	70
4.10	The amount of variance (%) of the first five principal components for two experiments	71
4.11	Mineral contents in raw unclean and clean EBN	76

© This item is protected by original copyright

LIST OF FIGURES

NO.		PAGE
2.1	Edible Bird Nest	7
2.2	World Range of Swiftlets	9
2.3	Cave Nest	12
2.4	House Nest	13
2.5	Structure of N-acetylneuraminic acid (sialic acid)	20
3.1	Summary of Experimental Procedures	29
3.2	EBN Solution	40
3.3	E-tongue setup for determination of nitrite and nitrate in EBN	41
4.1	FTIR spectra of raw unclean EBN samples (control), (a) Unclean Langkawi (b) Unclean Perlis (c) Unclean Java (d) Unclean Balikpapan (e) Unclean Kalimantan	46
4.2	FTIR spectra of clean EBN samples, (a) Clean Langkawi (b) Clean Perlis (c) Clean Java (d) Clean Balikpapan (e) Clean Kalimantan	47
4.3	FTIR spectra of samples, (a) EBN (b) Egg white (c) Sodium alginate (d) Carrageenan (e) Starch (f) Agar (g) Pork skin	49
4.4	FTIR spectrum of samples, (a) Pure EBN, (b) Starch + EBN, (c) Sodium alginate + EBN, (d) Carrageenan + EBN, (e) Agar + EBN, (f) egg white + EBN, (g) Pork skin + EBN	51
4.5	Moisture content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	53
4.6	Ash content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	55
4.7	Protein content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	57
4.8	Standard calibration curve for protein	57
4.9	Fat content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	60

4.10	Standard calibration curve for carbohydrate	61
4.11	Carbohydrate content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	62
4.12	Standard calibration curve for sialic acid	63
4.13	Sialic acid content of raw unclean and clean EBN from different geographical locations in Malaysia and Indonesia	64
4.14	Standard calibration curve for nitrite	66
4.15	Standard calibration curve for nitrate	67
4.16	Nitrite and nitrate content of raw unclean, clean and wastewater EBN from different geographical locations in Malaysia and Indonesia	67
4.17	PCA plot of nitrite content on raw unclean, clean and wastewater EBN samples from different geographical locations in Malaysia and Indonesia	72
4.18	PCA plot of nitrate content on raw unclean, clean and wastewater EBN samples from different geographical location in Malaysia and Indonesia	72

LIST OF SYMBOLS

$^{\circ}\text{C}$	Celsius
g	Gram
min	Minute
N	Normality
mg	Miligram
mg/L	Miligram per litre
ppm	Part per million
v/v	Volume to volume ratio
rpm	rotation per minute
α	Significance level
H_0	Hypothesis null
σ	Standard deviation

© This item is protected by original copyright

LIST OF ABBREVIATIONS

EBN	Edible Bird Nest
UL	Unclean Langkawi
UP	Unclean Perlis
UJ	Unclean Java
UB	Unclean Balikpapan
UK	Unclean Kalimantan
CL	Clean Langkawi
CP	Clean Perlis
CJ	Clean Java
CB	Clean Balikpapan
CK	Clean Kalimantan
ANOVA	Analysis of variance
Avg	Average
NO_2^-	Nitrite
NO_3^-	Nitrate
Na_2CO_3	Sodium carbonate
NaOH	Sodium hydroxide
CuSO_4	Copper sulphate
H_2SO_4	Sulphuric acid
HCl	Hydrochloric acid
H_2O_2	Hydrogen peroxide
HNO_3	Nitric acid

TFA	Trifluoroacetic acid
BSA	Bovine Serum Albumin
FTIR	Fourier Transform Infrared Spectroscopy
UV-Vis	Ultraviolet Visible Spectrophotometer
AAS	Atomic Absorption Spectroscopy
JMP software	JMP software from SAS Institute, USA (Version 10)
ISE	Ion selective sensor
E-tongue	Electronic tongue
AOAC	Association of Official Analytical Chemists
K	Potassium
Fe	Iron
Zn	Zinc
K	Potassium
Fe	Iron
Zn	Zinc
Cu	Copper
Pb	Lead

Komposisi kimia dan ketulenan sarang burung yang boleh dimakan terpilih daripada Malaysia dan Indonesia

ABSTRAK

Kajian ini dijalankan untuk menentukan keaslian EBN dan komposisi EBN yang bersih dan tidak bersih dan juga untuk membangunkan satu teknik pantas menggunakan teknologi sensor untuk menentukan kandungan nitrit dan nitrat. Sampel yang digunakan adalah EBN yang tidak bersih dan bersih daripada dua negara utama pengeksport: Malaysia (Langkawi dan Perlis) dan Indonesia (Java, Balikpapan dan Kalimantan) daripada spesies *Aerodramus fuciphagus* dan beberapa bahan tiruan iaitu agar, kanji, natrium alginat, rumpai laut, putih telur dan kulit babi. Pengenalpastian kumpulan berfungsi telah dikaji menggunakan teknik FTIR. Dalam kajian ini, kedua-dua EBN tulen yang tidak bersih dan bersih memberikan corak FTIR spektrum yang sama. Kehadiran kumpulan berfungsi seperti hidroksil (OH), karboksil (COOH), amina (NH) alkina (C-C), alkena (C=C), karbonil (C=O) dan nitro (N=O) telah dikenal pasti. Manakala, spektra bahan tiruan tulen adalah berbeza dengan EBN tulen. Spektra EBN tidak tulen dengan penambahan bahan tiruan menunjukkan persamaan dengan EBN tulen tetapi kehadiran kumpulan berfungsi adalah berbeza. Kawasan cap jari pada gelombang $<1500\text{ cm}^{-1}$ untuk spektrum EBN tulen dan EBN yang tidak tulen dengan penambahan bahan tiruan adalah berbeza dimana Kumpulan NH tidak hadir dalam semua bahan tiruan tulen kecuali kulit babi dan putih telur. Selain itu, hanya kulit babi mempunyai ikatan ester C=O berbanding EBN tulen dan bahan tiruan yang lain. Ikatan ester C=O ini adalah getaran daripada asid lemak pada gelombang 1745.1 cm^{-1} . Walaubagaimanapun, kedua-dua EBN dan bahan tiruan mempunyai ikatan C=O amide iaitu ikatan untuk protein pada gelombang $1670\text{-}1600\text{ cm}^{-1}$. Oleh itu, FTIR boleh digunakan sebagai kaedah yang mudah dan cepat untuk mengenalpasti ketulenan EBN dan bahan tidak halal. Komponen terbesar dalam EBN adalah protein dengan 22.4% hingga 32.7%. Kemudian, diikuti dengan kandungan karbohidrat dengan 9% hingga 12%. Kelembapan (6.92% hingga 13.88%), abu (1.1% hingga 8.8%) dan yang paling rendah adalah lemak mengandungi 0.01% hingga 0.07%. Nitrit dan nitrat dalam EBN telah ditentukan dengan menggunakan teknik sensor elektrod ion terpilih (ISE) juga dipanggil E-tongue. Teknik sensor adalah lebih cepat berbanding dengan teknik kromatografi ion biasa. Kepekatan nitrit dan nitrat dalam EBN tidak bersih, bersih dan air buangan EBN adalah berbeza dari 0.2 hingga 2.9 ppm, lebih rendah daripada bacaan standard iaitu 30 ppm dan selamat untuk dimakan oleh manusia. Ujian ketepatan yang telah dijalankan ke atas sarang burung yang boleh dimakan daripada Perlis menggunakan teknik sensor memberikan 100.2 hingga 100.8% ketepatan dengan nilai RSD kurang daripada 1%, manakala menggunakan teknik ion kromatografi memberikan 96.9 hingga 102.8% ketepatan dengan nilai RSD 9 hingga 13%. Justeru itu, teknik sensor mempunyai sensitiviti yang tinggi dan boleh digunakan sebagai teknik piawai dalam menentukan kandungan nitrit dan nitrat dalam sampel sarang burung yang boleh dimakan selain mengurangkan masa analisis daripada 2 jam kepada 10 minit. Unsur mineral seperti kalium dan besi adalah tinggi dalam EBN manakala kandungan plumbum adalah sangat rendah oleh itu ianya tidak beracun.

Chemical composition and authenticity of selected Edible Bird Nest in Malaysia and Indonesia

ABSTRACT

This study was conducted to determine the authenticity of EBN and the compositional properties of unclean raw and clean EBN as well as to develop a rapid technique to determine the nitrite and nitrate content using sensor technology. The samples used were raw unclean and clean EBN from two major exporting countries: Malaysia (Langkawi and Perlis) and Indonesia (Java, Balikpapan and Kalimantan) from species *Aerodramus fuciphagus* and some common adulterants namely agar, starch, sodium alginate, carrageenan, egg white and pork skin. Functional groups identification study was employed using FTIR technique. In this study, both unclean and clean EBN gave similar pattern of FTIR spectra. The presence of similar compounds such as hydroxyl (OH), carboxyl (COOH), amines (NH), alkanes (C-C), alkynes (C≡C), carbonyl (C=O) and nitro (N=O) groups were identified. The spectra of pure adulterant were apparently different with pure EBN while the spectra of adulterated EBN samples with addition of the adulterants were quite similar from that of pure EBN but the compounds present were different. The fingerprint region of the spectra of pure and adulterated edible bird nest samples were quite similar at wavenumber $<1500\text{ cm}^{-1}$. The NH group was absent in all the adulterants except in pork skin and egg white. Pork skin has ester C=O stretch bond which was not present in EBN itself and in other adulterants. The C=O stretch is an ester vibration band of fatty acids at wavenumber 1745.1 cm^{-1} . However, both EBN and adulterants have C=O stretch (amide) around $1670\text{-}1600\text{ cm}^{-1}$ for protein. Hence, FTIR can be used as a simple and rapid method to identify the authenticity of EBN and the non-halal component as adulterants. The largest component in EBN was protein with 22.4% to 32.7% respectively followed by carbohydrate ranging from 9% to 12%. Moisture (6.92% to 13.88%), ash (1.1% to 8.8%) and the lowest was fat content 0.01% to 0.07% respectively. Nitrite and nitrate in EBN was determined using sensor techniques using ISE electrode called E-tongue. This sensor technique is faster compared to the normal ion chromatography technique. Concentration of nitrite and nitrate in wastewater, unclean and clean EBN varies from 0.2 to 2.9 ppm which is lower than the standard set by China for export purposes. In this study, nitrite was less than 30 ppm and therefore safe for human consumption. The accuracy test carried out using Perlis EBN using sensor technique gave 100.2 to 100.8% recovery with RSD value less than 1% where as using ion chromatography gave 96.9 to 102.8% with RSD value 9 to 13%. Thus, sensor technique is highly sensitive and can be used as a standard method in determination of nitrite and nitrate in EBN samples as well as reduce the analysis time from 2 hour to 10 minutes. Mineral elements such as potassium and iron were high but lead content is low hence non-toxic.

CHAPTER 1

INTRODUCTION

1.1 Background of Study

Edible bird nest (EBN) refers to the nest produced by several different swiftlet species (Marcone, 2005). The common EBN traded around the world derived from two popular species which are white nest swiftlet (*Aerodramus fuciphagus*) and black nest swiftlet (*Aerodramus maximus*) (Babji et al., 2011). These nests were built from a glutinous material found in saliva secreted from the swiftlet's two sublingual salivary glands and may fuse with different materials like foliage and plumes (Chan, 2006).

EBN is generally mentioned as the 'Caviar of the East' as it is viewed as the most lavish creature nourishment item in the world. EBN from *Aerodramus* comprises of fat (0.14-1.28%), ash (2.1%), carbohydrate (25.62-27.26%) and protein (62.0-63.0%) (Marcone, 2005). EBN has been regarded as an esteemed food tonic for the Chinese people due to its highly evaluated function of being nutritious and medical benefit such as anti-aging, anti-cancer and immunity-enhancing. In Traditional Chinese Medicine (TCM), EBN has been utilized for a long time as a vital wellbeing supplement. EBN are also used for the treatment of malnutrition, a boost to the immune system and enhance body metabolism.

EBN comprises of high esteemed glycoprotein rich with amino acids, sugar, calcium, sodium and potassium (Norhayati et al., 2010). Sialic acid is the major component of carbohydrate (Colombo et al., 2003; Kathan and Weeks, 1969). Sialic acids have beneficial effect on neurological and scholarly focal points in babies (Chau et al., 2003). As

a superb insusceptible framework arbitrator, sialic acid influences the stream resistance in bodily fluid which thus repulses microorganisms, infections and other destructive organisms.

EBN is also rich in mineral salts where they contain high amount of sodium and calcium. It has been testified that the processed EBN contain calcium extended from 503.6 to 2071.3 mg/g and sodium content went from 39.8 to 509.6 mg/g which are higher than other mineral elements such as magnesium, potassium and others (Norhayati et al., 2010).

1.2 Statement of Problem

1.2.1 Authenticity of EBN

The high demand for EBN especially from China, Hong Kong and Taiwan and with limited supply has led to production of fake and adulterated EBN products in the global market (Set, 2012). The pure EBN has been adulterated with less expensive materials such as karaya gum, red seaweed, tremella fungus, sodium alginate, agar, natural plant gum, egg white and even the non-halal ingredient such as pork skin usually incorporated during the processing stages to increase the net weight or nutrient content prior to sale (Marcone, 2005).

Due to its high market value and indeed one of the most lavish animal food product in the world, the rate of adulteration of EBN is increasing (Koon and Cranbrook, 2002). Therefore, there is an urgent need to identify the authenticity of EBN. In this study, Fourier Transform Infrared Spectroscopy (FTIR) has been used to determine the authenticity of

EBN because it is a very simple and rapid technique. Another advantage is that no chemical pretreatment are used for the analysis.

1.2.2 Nitrite and Nitrate Content in EBN

The Malaysian EBN industry is facing problem of rejection from the importer especially China due to the high nitrite and nitrate content in EBN samples. The Chinese government demanded that nitrite must not be presented in EBN or its product in spite of the fact that the presence of natural nitrite in bird nest is being perceived. The presence of nitrite and nitrate in EBN have to comply with standards set by China and should not be more than 30 ppm (DSM, 2011).

In this study, the cleaning process of EBN is assessed for its effect to reduce nitrite and nitrate content in the wastewater, unclean and clean EBN. Isa (2012) reported that, cleaning process help to reduce nitrite and nitrate content in EBN by soaking them in the water. In conventional technique, the EBN was clean by soaking in the water until the nest is softened and then the small feathers and plumage were physically detached with tweezers.

1.3 Objective of the Study

The objectives of this study are:

1. To determine the authenticity of EBN from different locations in Malaysia and Indonesia.

2. To determine the compositional properties of raw unclean and clean EBN from different locations in Malaysia and Indonesia.
3. To develop a rapid technique to determine nitrite and nitrate in raw unclean, clean and wastewater EBN using sensor technology.

1.4 Scope of the study

This study was conducted to evaluate the quality of EBN that were collected from Malaysia: Perlis and Langkawi and Indonesia: Java, Balikpapan and Kalimantan. The authenticity of EBN were also determined to identify the pure EBN and faked or adulterated EBN. The compositional properties of EBN such as moisture, ash, protein, carbohydrate, fat, sialic acid and elemental contents were evaluated for quality, food value energy booster and health related properties. Since, one of the exportation parameter is nitrite and nitrate, the content of nitrite and nitrate in wastewater, unclean and clean EBN were also determined.

1.5 Chapter Summary

This chapter provided a brief background about the study, problem statement, objective and scope of the research. Edible bird nest which was constructed from the saliva of swiftlet consists of high amount of protein, carbohydrate and mineral which are good for human consumption as health supplement. However, the high demand of EBN has led to the production of fake EBN. So, there is an urgent need to identify the authenticity of EBN

using a rapid and environmentally friendly FTIR technique which is a rapid technique. The exportation of EBN to China has to comply with the standard set by China. The content of nitrite should not be more than 30 ppm. Therefore, a rapid technique using sensor was developed and evaluated for the determination of nitrite and nitrate in EBN.

© This item is protected by original copyright

CHAPTER 2

LITERATURE REVIEW

2.1 Edible Bird Nest (EBN)

Edible bird nest (EBN) is a dried glutinous secretion from the salivary glands of the male swiftlet in the breeding season (Guo et al., 2006; Marcone, 2005). It is also known as swiftlet nest produced by certain species of swiftlet family. During the nesting and breeding season, the male swiftlets construct their nest with glutinous strands of starched like saliva produce by a pair of large salivary gland under their tongue (Goh et al., 2000). The nests will increase its weight from 2.5 mg to 160 mg on reaching maximum secretory activity of swiftlets (Medway, 1962). The average young swiftlet will leave the nest after 90 to 120 days of weaning process and the nest will be collected (Tarburton, 2011).

As far as the nest constructing process in caves, swiftlets will use their saliva to bind materials together and attached them to the vertical walls of inland or seaside caves (Kang, Hails, & Sigurdsson, 1991). The nest hardened when exposed to air and looks like a cupped hand or half cut bowl about 3 to 5 inches in diameter. These nests are often referred as EBN (Figure 2.1) due to the fact that they supposedly contain a wide variety of nutrients for health and vitality , namely in relation to the lungs, libido and even skin complexion. Lately, it has been claimed as an anti-carcinogen (Chan, 2006).