



**Characterization and Optimization of Indoor
Environmental Quality on Grey Oyster (*Pleurotus
pulmonarius*) Mushroom Cultivation**

By

**Md. Tariqul Islam
(1433511528)**

A thesis submitted in fulfillment of the requirements for the degree of
Master of Science in Biosystem Engineering

**School of Bioprocess Engineering
UNIVERSITI MALAYSIA PERLIS**

2017

GRADUATE SCHOOL

UNIVERSITI MALAYSIA PERLIS

PERMISSION TO USE

In presenting this thesis in fulfillment of a post graduate degree from the Universiti Malaysia Perlis. I agree that permission for copying of this thesis in any manner, in whole or in part, for scholarly purposes may be granted by the Dean of the Graduate School. It is understood that any copying or publication or use of this thesis or parts thereof for financial gain shall not be allowed without any written permission. It is also understood that due recognition shall be given to me and to Universiti Malaysia Perlis for any scholarly use which may be made of any material from my thesis.

Request for permission to copy or to make other use of material in this thesis whole or in part should be addressed to

Dean, Centre for Graduate Studies

Universiti Malaysia Perlis (UniMAP)

112 & 114 (First Floor), Block A, Taman Pertiwi Indah,

Jalan Kangar – Alor Setar, Seriab

01000 Kangar, Perlis, Malaysia.

ACKNOWLEDGEMENT

In the name of Almighty ALLAH, The Most Gracious WHO blessed me to understand and carry out this work within my expected time-frame. I would like to present my greatest and sincerest thanks to HIM for gifting me knowledge and strength during the entire course of this Master durations.

I would like to honor my utmost gratitude to my main supervisor Assoc. Prof. Dr. Zarina Binti Zakaria, Faculty of Engineering Technology (FeTech), Universiti Malaysia Perlis (UniMAP), for her generous support, outstanding guidance, motivation, wonderful suggestions, discussion time allocated for me and strong faith upon me that she provided me throughout the work. Her continuous inspirations for working and publishing articles, constructive criticism and sincere willingness to solve problems have made this work to be completed within expected time. I would also like to thank my co-supervisor Assoc. Prof. Dr. Nasrul Bin Hamidin, Dean, School of Environmental Engineering, UniMAP, for his guidance and valuable suggestions during this research work. I heartily thank Assoc. Prof. Dr. Mohd Azlan Bin Mohd Ishak, Faculty of Applied Sciences, Universiti Teknologi MARA (UiTM), Malaysia, for outline during this research work. I would like to special thanks to UniMAP for providing me the financial support and others facilities to complete this research work.

I am grateful to all of my laboratory colleagues at School of Bioprocess Engineering (SBE) and FeTech, UniMAP, for their continuous support and friendly behaviors that encouraged me to work this research attentively. I would like to extend my sincere regards to postgraduate coordinator Dr. Saleha Binti Shamsuddin and all the

teaching and non-teaching staffs of School of Bioprocess Engineering for their co-operation and timely support. I am also grateful to Organic Mushroom Trading Bhd., Taman Bersatu, Arau, Perlis for giving consent during data collection in this research.

I am indebted to my dearest parents, elder brothers and sister, especially my lovely father (passed away January 15, 2014) Md. Abdul Wahab and mother Mrs. Amena Khatun who encourage me to achieve this goal. I am too much grateful to my elder brother Assoc. Prof. Dr. Asraf Ali, Daffodil International University, Bangladesh, who also awarded Ph.D. from School of Mechatronic Engineering, UniMAP, for his love, affection and generous supporting. Lastly, I am appreciative to my beloved wife Hosne Ara Sumi for her patience, substantial encouraging and mental supporting.

Thanks to Almighty ALLAH.

MD TARIQUL ISLAM

UNIVERSITI MALAYSIA PERLIS

TABLE OF CONTENTS

	PAGE
THESIS DECLARATION	i
PERMISSION TO USE	ii
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	v
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xv
ABSTRAK	xvii
ABSTRACT	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Background of Study	1
1.2 Problem Statement	6
1.3 Research Objectives	7
1.4 Research Scope	8
1.5 Thesis Organization	9
CHAPTER 2 LITERATURE REVIEW	10
2.1 Introduction to Mushroom	10
2.2 Mushroom Industry	11
2.2.1 Mushroom Production in the World	11
2.2.2 Mushroom Production in Malaysia	14
2.3 Nutritional and Medicinal Value	18
2.3.1 Nutritional Properties	19
2.3.2 Medicinal Properties	21

2.4	Mushroom Cultivation Substrate	22
2.5	Cultivation Methods	24
2.5.1	Cultivation on Logs	25
2.5.2	Cultivation on Beds	26
2.5.3	Cultivation on Sterilized Sawdust in Plastic Bags	26
2.5.4	Cultivation on Straw	28
2.6	Environmental Effects	28
2.6.1	Temperature	29
2.6.2	Relative Humidity (RH)	32
2.6.3	Carbon Dioxide (CO ₂)	35
2.6.4	Light	36
2.7	Indoor and Outdoor Environment of Malaysia	38
2.8	Mushroom Cultivation in Controlled Environment	42
	CHAPTER 3 RESEARCH METHODOLOGY	44
3.1	Preparation of Mushroom Growing Bags	44
3.2	Model and Design of Indoor Cultivation Room	47
3.3	Experimental Design and Controlled Mushroom Cultivation	49
3.3.1	Characterization of Indoor Environment	49
3.3.1.1	Ambient Indoor Environment Without Humidifying and Ventilation (System 1)	51
3.3.1.2	Indoor Humidifying Environment Without Ventilation (System 2)	51
3.3.1.3	Indoor Humidifying Environment With Ventilation (System 3)	53
3.3.1.4	Cultivation in Humidifying Till Primordial Initiation Without Ventilation (System 4)	54
3.3.1.5	Cultivation in Humidifying Till Primordial Initiation With Ventilation (System 5)	54

3.3.1.6	Data Collection	55
3.3.1.6.1	Environment Conditions	55
3.3.1.6.2	Morphology and Moisture Contents	56
3.3.1.6.3	Mushroom Yield	57
3.3.1.7	Statistical Analysis	58
3.3.2	Optimization of Indoor Environment	58
3.3.2.1	Indoor Humidifying Environment With Ventilation (System 6)	61
3.3.2.1.1	Combined Application	62
3.3.2.1.2	Individual Application	63
	Configuration 1	63
	Configuration 2	64
	Configuration 3	66
	Configuration 4	66
	Configuration 5	68
3.3.2.1.3	Cultivation in Optimized Procedure	68
3.3.2.2	Cultivation in Outdoor Environment (System 7)	69
3.3.2.3	Data Collection	71
3.3.2.3.1	Environment Conditions	71
3.3.2.3.2	Morphology and Moisture Contents	71
3.3.2.3.3	Total Yield	72
3.3.2.4	Statistical Analysis	72
3.4	Biochemical Analysis	72
3.4.1	Preparation of sample	73
3.4.2	Determination of Total Protein	74
3.4.2.1	Preparation of Protein Standard Curve	74

3.4.2.2	Total Protein Analysis	74
3.4.3	Determination of Total Carbohydrate	75
3.4.3.1	Preparation of Glucose Standard Curve	75
3.4.3.2	Total Carbohydrate Analysis	76
3.4.4	1) Determination of Total Fat	77
CHAPTER 4 RESULTS AND DISCUSSION		78
4.1	Characterization of Indoor Environment	78
4.1.1	Optimize Humidifying Duration	78
4.1.1.1	System 3	78
4.1.1.2	System 4	80
4.1.2	Environmental Conditions	82
4.1.3	Mushrooms Morphology and Moisture Contents	88
4.1.3.1	Individual Weight (IW)	88
4.1.3.2	Numbers of Fruitbodies (NFB)	90
4.1.3.3	Cap Diameter (CD)	92
4.1.3.4	Stalk Height (SH)	92
4.1.3.5	Moisture Contents	94
4.1.4	Mushroom Yield	98
4.1.4.1	Primordial Initiation	98
4.1.4.2	Mature Fruitbodies Formation	99
4.1.4.3	Dead Primordia Formation	100
4.1.4.4	Bio-efficiency	101
4.2	Optimization of Indoor Environment	103
4.2.1	Optimize Humidifying Duration	103
4.2.2	Optimized Humidifier Configuration	104

4.2.3	Environmental Conditions	108
4.2.3.1	Hourly Temperature and Humidity	108
4.2.3.2	Overall Temperature	110
4.2.3.3	Overall Relative Humidity	110
4.2.3.4	Carbon Dioxide (CO ₂)	112
4.2.3.5	Light Intensity	113
4.2.4	Morphology and Moisture Contents	113
4.2.5	Mushroom Yield	122
4.2.5.1	Total Yield	122
4.2.5.2	Bio-efficiency	124
4.3	Correlation Among Environmental and Morphological Parameters	125
4.4	Biochemical Analysis	127
4.3.1	Protein Analysis	127
4.3.1.1	Protein standard Calibration Curve Analysis	127
4.3.1.2	Protein Content in Different Cultivation Systems	128
4.3.2	Total Carbohydrate Analysis	130
4.3.2.1	Standard Curve Calibration Analysis	130
4.3.2.2	Carbohydrate Content in Different Cultivation Systems	130
4.3.3	Lipid Analysis	132
CHAPTER 5 CONCLUSIONS AND RECOMMENDATIONS		136
5.1	Finding and Contributions	136
5.2	Recommendation for Future Works	138
REFERENCES		140
APPENDIX A	Specifications of Materials	150
APPENDIX B	Bovine Serum Albumin (BSA) Standard Curve	153

APPENDIX C	Sample Protein Analysis	157
APPENDIX D	Standard Glucose Curve and Sample Carbohydrate Analysis	163
APPENDIX E	Sample Lipid Analysis	168
APPENDIX F	Gallery	170
LIST OF PUBLICATIONS AND AWARDS		178

©This item is protected by original copyright

LIST OF TABLES

NO		PAGE
2.1	World production of mushrooms, 1990-2012 (tonnes)	13
2.2	Types of mushrooms cultivated in Malaysia	15
2.3	Number of mushroom growers and mushroom production in Malaysia.	16
2.4	Import and export of mushroom in Malaysia 2001-2007.	17
2.5	Nutritional contents of different oyster mushrooms (g/100g).	20
2.6	Nutritional contents of <i>Pleurotus pulmonarius</i> in different studies.	21
2.7	Optimal temperature ($^{\circ}\text{C}$) of different oyster mushrooms in different stages of growing.	31
2.8	Optimal humidity (%) different oyster mushrooms in different stages of growing.	34
3.1	Experimental design of controlled mushroom cultivation for characterization of indoor environment	49
3.2	Sample collection from different cultivation systems for biochemical analysis.	73
4.1	Calculation of optimized humidifying duration for System 2 and System 3.	81
4.2	Descriptive statistics of temperature and humidity in different indoor mushroom cultivation systems.	83
4.3	Least Significant Difference (LSD) analysis of temperature and humidity for characterization of indoor environment. The mean difference is significant at $p < 0.05$.	85
4.4	Descriptive statistics of mushroom morphology for characterization of indoor environment.	89
4.5	One-way ANOVA Analysis for the morphology of <i>P. pulmonarius</i> among the cultivation systems.	91
4.6	LSD Analysis of mushroom's morphology in different cultivation systems for characterization.	93
4.7	Descriptive of humidity and temperature in different ways of humidifier configuration.	106

4.8	Overview of environmental factors (temperature, humidity, CO ₂ and light) during the cultivation of System 6 and System 7.	111
4.9	One-way ANOVA Analysis of environmental factors during the cultivation of <i>Pleurotus pulmonarius</i> in System 6 and System 7.	112
4.10	One-way ANOVA analysis of Mushrooms morphology and quality in System 6 and System 7.	119
4.11	Correlation among with the environmental factors and the morphology of <i>P. pulmonarius</i> .	126
4.12	Percentages of protein concentration in different cultivation systems (Dried Sample).	128
4.13	One-way ANOVA analysis for protein contents of mushroom in different cultivation Systems.	129
4.14	Percentages of carbohydrate concentration in different cultivation system (Dried sample).	131
4.15	One-way ANOVA analysis for carbohydrate contents of mushroom in different cultivation systems.	132
4.16	Percentages of lipid concentration in different cultivation systems (g/100g Dried Sample).	133
4.17	One-way ANOVA analysis for lipid contents of mushroom in different cultivation systems.	134

LIST OF FIGURES

NO		PAGE
1.1	Geographic Map of Malaysia with latitude and longitude direction presenting the temperature in April 2016.	4
2.1	Morphological structure of mushroom.	11
2.2	Different methods for mushroom cultivation.	27
2.3	The average indoor relative humidity and air temperature all over the year in Malaysia (8:00 am to 12:00 am).	39
2.4	The average outdoor relative humidity and air temperature all over the year in Malaysia (8:00 am to 12:00 am).	40
2.5	Daily air temperature and humidity patterns for daytime ventilation, night ventilation and outdoor in Malaysia.	41
3.1	Steps of mushroom growing bags preparation	45
3.2	Mycelium colonization; 0-5% (A); 40-50% (B); 70% (C) and 100% (D) completed.	45
3.3	Flow diagram of research methodology	46
3.4	Floor plan and design of indoor mushroom cultivation house with bags	48
3.5	Flowchart of the protocol for the characterization of indoor environment during mushroom cultivation.	50
3.6	Arrangement of bags and humidifier for characterization of indoor environment.	52
3.7	Flowchart of the protocol for the optimization of indoor environment during mushroom cultivation.	60
3.8	Configuration of humidifier position in different points inside the growing house.	62
3.9	Humidifying configuration 1 for individual application	64
3.10	Humidifying configuration 2 for individual application	65
3.11	Humidifying configuration 3 for individual application	65
3.12	Humidifying configuration 4 for individual application.	67
3.13	Humidifying configuration 5 for individual application.	67

3.14	Side view of indoor mushroom cultivation in optimized condition of System 6.	69
3.15	Side view of outdoor mushroom growing farm (System 7) without black net and bags (left), with surrounding black net and bags (Right).	70
4.1	Optimization of humidifying duration in indoor environment without ventilation system (System 2). Values are mean of 4 replicates.	79
4.2	Optimization of humidifying duration in indoor environment with ventilation system (System 3). Values are mean of 4 replicates.	80
4.3	Percentages of moisture contents of <i>P. pulmonarius</i> fruitbodies in different cultivation system.	95
4.4	The percentages of primordia initiation, mature fruitbody formation and dead primordia formation in different environmental cultivation system.	100
4.5	Variation of bio-efficiency (BE %) in the cultivation system of different environmental conditions.	102
4.6	Graph for increased humidity in indoor environment for System 6. Values are mean of 4 replicates.	104
4.7	Hourly variation of mean temperature and humidity in the cultivation of System 6.	108
4.8	Hourly variation of mean temperature and humidity in the cultivation of System 7.	109
4.9	Mean difference of individual weight by numbers of cycle in the cultivation of System 6 and System 7.	114
4.10	Mean difference of numbers of fruitbodies by numbers of cycle in the cultivation of System 6 and System 7.	115
4.11	Mean difference of cap diameter by numbers of cycle in the cultivation of System 6 and System 7.	117
4.12	Mean difference of stalk height by numbers of cycle in the cultivation of System 6 and System 7.	118
4.13	Percentages of moisture containing fruitbodies in different number of cycles in the cultivation of System 6.	120
4.14	Percentages of moisture containing fruitbodies in different number of cycles in the cultivation of System 7.	121
4.15	Total yield performance in the cultivation of System 6 and System 7.	123

LIST OF ABBREVIATIONS

RH	Relative Humidity
MC	Moisture Contents
TWFFB	Total weight of Fresh Fruitbodies
DWFB	Dry Weight of Fruitbodies
PPI	Percentages of Primordial Initiation
NBPI	Number of Bags of Primordial Initiation
TNB	Total Numbers of Bag
PMF	Percentages of Mature Fruitbodies
NBMF	Number of Bags of Mature Fruitbodies
PDP	Percentages of Dead Primordia
BE	Bio-Efficiency
TWFB	Total Weight of Fruitbodies from each Bags
TWS	Total Weight of Substrate of each Bags
H1	Humidifier 1
H2	Humidifier 2
H3	Humidifier 3
H4	Humidifier 4
I	Interval
Kg	Kilogram
Cm	Centimetre
M	Meter
BSA	Bovine Serum Albumin
ML	Millilitre

DF	During Fruiting
DPI	During Primordial Initiation
IW	Individual Weight
NFB	Numbers of Fruitbodies
CD	Cap Diameter
SH	Stalk Height
MD	Mean Difference
LSD	Least Significant Difference
ANOVA	Analysis of Variance
MOA	Ministry of Agriculture Malaysia
<i>P.</i>	<i>Pleurotus</i>
<i>spp.</i>	<i>Species</i>
FAO	Food and Agriculture Organization
Min	Minute
RPM	Rotation per minute
L	Litre

Pencirian dan Pengoptimuman Kualiti Persekitaran Dalaman Terhadap Penanaman Cendawan Tiram Kelabu (*Pleurotus pulmonarius*)

ABSTRAK

Berdasarkan pemerhatian terhadap peningkatan permintaan cendawan, pengusaha cendawan menghadapi rintangan yang lebih mencabar disebabkan perubahan keadaan persekitaran yang tidak menentu. Mengawal dan memantau kualiti persekitaran di dalam penanaman cendawan adalah faktor penting untuk pengeluaran cendawan yang tinggi. Oleh itu, kajian ini dilakukan untuk memeriksa kesesuaian dan perkembangan pertumbuhan *P.pulmonarius* di dalam persekitaran dalaman. Untuk ini, rekabentuk kajian dijalankan dengan dua prosedur berbeza iaitu pencirian dan pengoptimuman persekitaran dalaman. Pencirian dijalankan dengan lima keadaan persekitaran yang berbeza iaitu Sistem 1 sehingga Sistem 5. Substrat untuk cendawan berlambang telah disediakan dengan mencampurkan habut kayu, dedak dan kapur pertanian untuk semua sistem penanaman dan behih telah disuntik. Dengan menggunakan substrat yang dipenuhi miselium, eksperimen tanaman cendawan telah dijalankan menggunakan Sistem 1 sehingga Sistem 5. Sistem 1 adalah persekitaran dalaman secara semula jadi manakala Sistem 2 hingga Sistem 5 menggunakan kaedah persekitaran dalaman berkelembapan dengan atau tanpa pengudaraan. Purata suhu dalaman paling rendah dan kelembapan tertinggi adalah pada Sistem 3 yang mana menunjukkan perbezaan ketara dengan semua sistem yang lain. Morfologi tertinggi dan kandungan kelembapan jana buah luar biasa juga ditemui di Sistem 3. Morfologi keseluruhan dan kandungan kelembapan jana buah di Sistem 3 telah menunjukkan perbezaan yang ketara berbanding dengan hasil dari sistem lain. Sebanyak 98.4% permulaan primodia dan 98.6% jana buah matang dengan 1.4% primodia mati ditemui di Sistem 3. Bacaan primodial terendah iaitu sebanyak 30.2% ditemui di Sistem 1. Bacaan terendah jana buah sebanyak 43.1% dan 42.4 %, dan bacaan tertinggi primodia mati adalah 56.9% dan 57.6% masing-masing ditemui di Sistem 4 dan Sistem 5. Bio-kecekapan (73.6%) tertinggi telah dicapai dalam penanaman Sistem 3. Prosedur pengoptimuman telah dijalankan oleh dua keadaan persekitaran berbeza iaitu Sistem 6 dan 7. Sistem 6 dijalankan seperti Sistem 3 dengan menambah jumlah alat kelembapan, pengudaraan dan jumlah beg substrat. Untuk ini, alat kelembapan dan pengudaraan nafas telah digunakan dengan satu gabungan dan lima aplikasi individu untuk mengoptimumkan tempoh kelembapan dan prosedur penanaman. Manakala Sistem 7 menjalankan penanaman secara luaran untuk perbandingan hasil kajian. Konfigurasi optima ditemui sebagai aplikasi selama 15 minit rawatan kelembapan dengan diikuti selang masa selama 15 minit bagi setiap alat kelembapan. Suhu yang paling rendah dan kelembapan paling tinggi didapati dalam Sistem 6. Morfologi dan kandungan kelembapan jana buah didapati paling tinggi di Sistem 6, dimana jumlah hasil sebanyak 261.8 kg yang mana 140.3 kg lebih tinggi daripada Sistem 7. Bacaan suhu menunjukkan hubungan negatif yang ketara dan kelembapan menunjukkan hubungan positif yang ketara dengan morfologi dan kandungan kelembapan di dalam jana buah. Sebaliknya, kandungan kelembapan, bilangan jana buah, diameter tepi jana buah dan ketinggian tangkai telah menunjukkan hubungan positif yang ketara kepada berat setiap jana buah. Protein, karbohidrat, dan kandungan lipid didapati sama seperti dalam keadaan penanaman biasa dan tidak ada perbezaan yang signifikan didapati dalam sistem penanaman. Hasil kajian ini, memainkan peranan utama untuk penanaman cendawan tertutup dengan menyediakan garis panduan tertentu untuk mengawal faktor-faktor persekitaran, pembangunan sistem penanaman yang lebih baik, meningkatkan morfologi cendawan, kualiti dan hasil, risiko rendah pencemaran dan akhir sekali bagi memastikan pengeluaran berterusan cendawan bebas dari cuaca untuk sepanjang tahun.

Characterization and Optimization of Indoor Environmental Quality on Grey Oyster (*Pleurotus pulmonarius*) Mushroom Cultivation

ABSTRACT

In view of increasing demand, mushroom cultivators are facing more challenges and difficulties to grow mushroom due to inconsistent environmental conditions. Controlling and monitoring of environmental quality in mushroom cultivation is an important factor for high mushroom production. Therefore, this study was initiated to examine the suitability and growing performance of *P. pulmonarius* in indoor environmental condition. For this, the experimental design was followed by two different procedures as characterization and optimization of the indoor environment. Characterization procedure was conducted by mushroom cultivation in five different environmental conditions including from System 1 to System 5. The substrate for mushroom growing was prepared by mixing of sawdust, rice bran and agricultural lime for the all cultivation systems and spawn was inoculated. After completing mycelium colonization the experimental mushroom cultivations were conducted by System 1 to System 5. System 1 was the natural indoor environment whereas from System 2 to System 5 were the indoor humidifying environment with or without ventilation. The lowest mean indoor temperature and the highest mean humidity was found in System 3 which showed significant results with all other systems. The highest morphology and outstanding moisture containing fruitbodies were also found in System 3. The highest 98.4% primordial initiation and 98.6% mature fruitbodies formation with the lowest 1.4% dead primordia were found in System 3. The lowest 30.2% primordial initiation was found in System 1. The lowest 43.1% and 42.4% mature fruitbodies and the highest 56.9% and 57.6% dead primordia were found in System 4 and 5 respectively. The highest bio-efficiency (73.6%) of yield was achieved in the cultivation of System 3. The optimization procedure was conducted under two different environmental conditions as named System 6 and System 7. System 6 was followed by the procedure of System 3 with additional numbers of the humidifier, ventilation, and substrate bags. For this, the humidifiers and ventilations were applied by one combined and five individual treatments to optimize the humidifying duration and cultivation procedure. On the other hand, outdoor cultivation was conducted as System 7 for comparison study. The optimized configuration was found as the application of 15 mins humidifying treatment and followed by 15 mins interval period for each humidifier. The lowest temperature and highest humidity were found in System 6. The morphology and moisture contents of fruitbodies were found significantly high in System 6, where the total yield was found 261.8 kg which was 140.3 kg higher than System 7. The temperature showed significant negative correlation and the humidity showed significant positive correlation with the morphology and moisture contents of fruitbodies. On the other hand, moisture contents, numbers of fruitbodies, cap diameter and stalk height had shown significant positive correlation with individual weight. The protein, carbohydrate, and lipid contents were found similar to conventional cultivation and no significant different was found among the cultivation systems. The outcomes of the present research may play a key role in indoor mushroom cultivation by specific guideline to control environmental factors, develop better cultivation system, improve mushrooms morphology, quality, and yield, low risk of contamination and finally ensure continuous weather independent mushroom production throughout the years.

CHAPTER 1

INTRODUCTION

1.1 Background of Study

Mushroom cultivation in Malaysia is getting important due to its outstanding nutritional and medicinal properties which are 100% vegetarian food and is good for the patient of diabetes and joint pains. The demand of mushrooms especially the grey oyster mushroom (*Pleurotus pulmonarius*) is greater than their production. It's easy to consume and offers high income to local growers due to its ability to grow and fruiting in subtropical and tropical region. *P. pulmonarius* also known as *P. sajor-caju*, is a valuable edible mushroom, commonly called the grey oyster mushroom (Avin, 2014).

In Malaysia, the mushroom has been cultivated since the early 1970s and now mushroom is declared as an industrial crop. As an industrial crop, the mushroom has been included in National Agro-Food Policy Malaysia 2011-2020. The objective of this policy is to ensure sufficient mushroom supplies and to increase the income of cultivators. The most strategies of this policy are to increase production and productivity and control the quality of production. In the policy, the estimated target of mushroom production has to increase by 16 % in every year, from 15 thousand metric tons in 2010 to 67 thousand metric tons in 2020. Where the export value projected to increase about RM300 million in 2020 at the rate of 16.1 % per year (Ministry of Agriculture Malaysia, 2011).

Typically, the number of mushroom growers in different states of Malaysia are fluctuating, because of unstable production and market changes. However, the total

number of growers in Malaysia has reportedly increased in every year. In 2007, the total cultivators were 339 which are reached 428 in 2014. Johor, Selangor, and Pahang states are leading in mushroom producers, probably because of the higher demand especially from Chinese Malaysian population (Rosmiza, Davies, Jabil, & Mazdi, 2016). But most of them (about 80%) are small growers and produce below 50 kg per hectare of fresh mushrooms in a day. The medium scale (produce 50-500 kg per day) growers constitute 17% and big scale industries (producing more than 500kg per day) are around 3% only (Haimid, Rahim, & Dardak, 2013).

Although the numbers of mushroom growers in Malaysia are increasing but a lot of mushrooms have to import into Malaysia from other countries mainly from China, Thailand, Hong Kong and Taiwan which is still increasing. According to the report of Rosmiza et al. (2016), mushroom export has increased by about 19% in a year from RM12 million in 2000 to RM67 million in 2008. But the exports slightly dropped from 8302.96 metric tons in 2008 to 3190.26 metric tons in 2013. On the other hand, the import has increased from more than 21 thousand tons in 2004 to 24.5 thousand tons in 2013. This increment of import being partly due to less production, increasing population and a reported higher concern towards health (Mohd-Syauqi, Tapsir, Alam, Hasnul, & Mohd-Zaffrie, 2014).

Together with increasing numbers of mushroom growers, they are not able to produce more mushroom as demand. According to the Ministry of Agriculture 2010 (Haimid et al., 2013), in local market daily demand for fresh mushroom is 50,000 kg while the supply is only 24,000 kg. The most important fact is that the mushroom growers are facing more challenges and difficulties to grow mushrooms due to the inconsistent environmental conditions. The environment is considering the most important physical factor for mushroom production and quality. Referring to the

Agriculture and Food Development Authority Malaysia (2013), the production deficit of mushroom in the year 2009 was approximately 11,000 tons than the previous year. This deficit happened mainly due to the inconsistent environmental conditions of Malaysia by high temperature and low humidity.

This reduction of mushroom production is mainly due to the variation of environmental conditions because most of the mushroom cultivators depend on the outdoor cultivation. So, they are unable to control the inconsistent weather of high temperature and low humidity. They just depend on the natural weather of Malaysia that can reduce the production, morphology, and quality. Even, sometimes this type of mushroom have more risk for pest infection that can be an unhygienic food to the consumers (Fletcher & Gaze, 2007; Singh & Sharma, 2016).

The geography of Malaysia deals with the tropical climate, a country located in Southeast Asia. Malaysia is situated in the equatorial doldrums' area, has a geographic coordinate that reads 2° 30' North latitude and 112° 30' East longitude. Malaysia has tropical weather, influenced by monsoonal climate because of its latitude and longitude (Figure 1.1). Tropical climate here gives hot summer that is accompanied with low humidity level. So, the weather of Malaysia is generally hot and sunny throughout the year especially during February to October. During the hot season, the temperature may rise up to 41°C which may cause very low relative humidity.




Figure 1.1: Geographic Map of Malaysia with latitude and longitude direction presenting the temperature in April 2016 (Malaysian Metrological Department, 2016).

In the annual report of Malaysian Metrological Department (2015), the maximum 31.2 to 34.1 °C and minimum 22.3 to 24.4 °C temperatures was varied throughout the year whereas the maximum relative humidity at day was varied 60-67% and 94-96% at night. Moreover, the temperature and humidity in both indoor and outdoor environment are shown inversely proportional to each other. The indoor humidity is lower than outdoor at night since at night the indoor temperature was found higher than the outdoor environment (Wafi, Ismail, & Ahmed, 2011). The highest outdoor temperature of this running year 2016 was recorded 39.3 °C in April which caused the lowest relative humidity (Malaysian Metrological Department 2016, Figure 1.1). Whereby, *Pleurotus* species able to grow in 25-30 °C temperature and 80-90% humidity (Newsham, 2012; Sher, Al-Yemini, Bahkali, & Sher, 2010; Uddin et al., 2011).

So, the environment seems to be very important factors for induction of primordial initiation and fruitbodies formation. The appropriate environment is not only crucial for production, but also influences the ideal morphology and quality. Inappropriate temperature and humidity inhibit the mushroom growth and can even wilt and die. Most studies focusing on the relationship of mushroom cultivation with spawn, substrate and nutritional value. Some studies had performed on the environment to find out suitable species for the suitable season in the multi-seasonal country. Some studies also investigated the mushroom production in the artificially controlled indoor environment to find out the optimum ranges of humidity, temperature, carbon-dioxide and light intensity for maximum bio-efficiency (Jang et al., 2003; Seung, Byun, You, & Park, 1984; Yang et al., 2013). For controlling the environment, most of the studies used the air conditioner, evaporative cooler, growth chamber and other different cooling systems which are expensive, high operation costing and not suitable for large or commercial cultivation.

As the temperature and humidity of Malaysia weather is far from the optimal condition for oyster mushroom cultivation. With the traditional or conventional method used by most cultivators, is difficult to achieve the maximum yield as required. Moreover, it is important to know the characteristics of indoor environment for mushroom cultivation in Malaysia. Therefore, this study was conducted to develop a design and optimization procedure to control the indoor environment for the maximum growing performance that leads to ensure continuous mushroom production throughout the year independence of weather and climate change. Moreover, this study is very much relevance to the 2011-2020 Malaysia Agro-food Policy which one of the objectives is to enhance mushroom production.

1.2 Problem Statement

Many studies showed that oyster mushroom can grow ranging between 25 °C to 30 °C of temperatures and 80% to 90% of relative humidity (Miles & Chang, 2004; Uddin et al., 2011). According to the official website of Malaysian Meteorological Department (MMD), weather of Malaysia is generally hot and sunny all over the year especially from February to October. The temperature can rise up to 41 °C which can cause low relative humidity by 58% during the hot season. When temperature value reaches 38 °C or above, the mushroom production may drop by 50% and the quality also low as mushroom will dry, deformed, hard fruitbodies, which are unfavorable to the customers (Cha, 2004; Jang et al., 2003).

Generally, the cultivators have used a sprinkler or watering system which not able to maintain favorable humidity and reduce temperature due to the open environment. An open system increases the loss of humidity and therefore increases the temperature even with the frequent watering application. A closed cultivation system may help to reduce the loss of humidity from the environment and able to control by optimal ranges. For the time being, no study is conducted on the characterization and performance of indoor mushroom cultivation. Some research works were had conducted to grow mushroom production in artificially controlled environment to investigate the optimum requirements for maximum bio-efficiency (Jang et al., 2003; Seung et al., 1984; Yang et al., 2013) by using air conditioner, evaporative cooler, growth chamber and other different cooling systems which are expensive, high operation costing and not suitable for large or commercial cultivation. So, an optimized procedure should be investigated to control the optimal humidity (80-90%) and tolerable temperature (lower